

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 July 2001 (12.07.2001)

PCT

(10) International Publication Number
WO 01/49682 A1

(51) International Patent Classification⁷: C07D 403/12, 495/04, 471/04, 487/04, 498/04, A61K 31/55, A61P 13/00 // (C07D 495/04, 223:00, 333:00) (C07D 495/04, 243:00, 333:00) (C07D 471/04, 221:00, 243:00) (C07D 487/04, 231:00, 243:00) (C07D 487/04, 209:00, 255:00) (C07D 487/04, 243:00, 249:00) (C07D 487/04, 209:00, 243:00)

Baddesley, Southampton, Hampshire SO52 9NQ (GB). YEA, Christopher, Martyn [GB/GB]; 11 Gaston Gardens, Romsey, Hampshire SO51 7RZ (GB). FRANKLIN, Richard, Jeremy [GB/GB]; 208 Finchampstead Road, Wokingham, Berkshire RG40 3HB (GB).

(21) International Application Number: PCT/GB01/00023

(74) Agent: GEERING, Keith, Edwin; Reddie & Grose, 16 Theobalds Road, London WC1X 8PL (GB).

(22) International Filing Date: 4 January 2001 (04.01.2001)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0000079.4 5 January 2000 (05.01.2000) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): FERRING BV [NL/NL]; Polaris Avenue 144, NL-2132 JX Hoofddorp (NL).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): ASHWORTH, Doreen, Mary [GB/GB]; 10 The Glades, Locks Heath, Southampton, Hampshire SO31 6UY (GB). PITT, Gary, Robert, William [GB/GB]; 18 Kingfishers, Ship-ton-Bellinger, Tidworth, Hampshire SP9 7US (GB). HUDSON, Peter [GB/GB]; 12 Totteahale Close, North


Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CONDENSED AZEPINES AS VASOPRESSIN AGONISTS

WO 01/49682 A1



(57) Abstract: This invention provides novel compounds according to general formula (1) wherein A is a bicyclic or tricyclic azepine derivative, V¹ and V² are both H, OMe or F, or one of V¹ and V² is Br, Cl, F, OH, OMe, OBn, OPh, O-acyl, N₃, NH₂, NHBn or NH-acyl and the other is H, or V¹ and V² together are =O, -O(CH₂)_pO- or -S(CH₂)_pS-; W¹ is either O or S; X¹ and X² are both H, or together are =O or =S; Y is OR⁵ or NR⁶R⁷; R¹, R², R³ and R₄ are independently selected from H, lower alkyl, lower alkyloxy, F, Cl and Br; R⁵ is selected from H and lower alkyl; R⁶ and R⁷ are independently selected from H and lower alkyl, or together are -(CH₂)_n; n=3, 4, 5, 6; and p is

2 or 3. The compounds are agonists at the vasopressin V₂ receptor and are useful as antidiuretics and procoagulants. The invention further comprises pharmaceutical compositions incorporating these vasopressin agonists, which compositions are particularly useful in the treatment of central diabetes insipidus, nocturnal enuresis and nocturia.


CONDENSED AZEPINES AS VASOPRESSIN AGONISTS

FIELD OF INVENTION


The present invention relates to a class of novel chemical entities which act as agonists of the peptide hormone vasopressin. They reduce urine output from the kidneys and so are useful in the treatment of certain human diseases characterised by polyuria. They are also useful in the control of urinary incontinence and bleeding disorders.

BACKGROUND TO THE INVENTION

Vasopressin is a peptide hormone secreted by the posterior pituitary gland. It acts on the kidney to increase water retention and so reduce urine output. For this reason, vasopressin is alternatively known as "antidiuretic hormone". It also acts on the vasculature, where it produces a hypertensive effect. The cellular receptors that mediate these two actions have been characterised and shown to be different. The antidiuretic action is mediated by the type-2 vasopressin receptor, commonly called the V₂ receptor. Agents that can interact with the V₂ receptor and activate it in the same way as vasopressin are called V₂ receptor agonists (or simply V₂ agonists). Such agents will have an antidiuretic action. If these agents interact selectively with the V₂ receptor and not the other vasopressin receptor subtypes, then they will not have the hypertensive effect of vasopressin. This would be an important safety consideration and make such agents attractive for the treatment of human disease conditions characterised by polyuria (which is herein taken to mean excessive urine production).



In fact, such an agent is already in use in human therapy. Desmopressin (otherwise [1-desamino, D-Arg⁸]vasopressin, MinirinTM, DDAVPTM) is a peptide analogue of vasopressin which is selectively an agonist at the V₂ receptor. It is used in the treatment of central diabetes insipidus, which is a condition that results from defective secretion of vasopressin. It is also employed in the control of nocturnal enuresis and may also be of use in the control of nocturia. However, desmopressin is not an ideal agent in all respects. Even the best current syntheses of the agent are lengthy, and desmopressin is not amenable to the most convenient of purification techniques such as crystallisation. Consequently, desmopressin is relatively expensive. It has a very low oral bioavailability, and there is some variability in this parameter.




Overall then, there exists a need for a selective vasopressin V₂ receptor agonist that is easy to prepare and purify, and that has a high and predictable oral bioavailability. Such properties are most likely to be obtained with a non-peptide compound. These considerations have led other groups to investigate non-peptide vasopressin V₂ agonists, and their results are disclosed in, for example, International Patent Applications WO97/22591, WO99/06403, WO99/06409, WO00/46224, WO00/46225, WO00/46227 and WO00/46228. The compounds disclosed in these documents are, however, less than ideal. In particular, they have poor oral bioavailability, probably due in part to their low aqueous solubility. The present invention provides compounds with improved solubility and bioavailability.

Besides its antidiuretic actions, desmopressin is used to increase the concentration in the blood of the coagulation proteins known as Factor VIII and von Willebrand factor. In the clinical context, this makes desmopressin useful in the treatment of haemophilia A and von Willebrand's disease. Similar applications would be open to the non-peptide agonists of the present invention.







SUMMARY OF THE INVENTION

As disclosed herein, the present invention relates to a series of compounds that are non-peptide agonists of vasopressin and which are selective for the V_2 receptor subtype. The compounds are described by general formula 1



wherein:

A is a bicyclic or tricyclic azepine derivative selected from general formulae 2 to 7



A^1 , A^4 , A^7 and A^{10} are each independently selected from CH_2 , O and NR^8 ;

A^2 , A^3 , A^9 , A^{11} , A^{13} , A^{14} and A^{15} are each independently selected from CH and N;

either A^5 is a covalent bond and A^6 is S, or A^5 is $N=CH$ and A^6 is a covalent bond;

A^8 and A^{12} are each independently selected from NH and S;

A^{16} and A^{17} are both CH_2 , or one of A^{16} and A^{17} is CH_2 and the other is selected from O, SO_x , and NR^8 ;

V^1 and V^2 are both H, OMe or F, or one of V^1 and V^2 is Br, Cl, F, OH, OMe, OBn, OPh, O-acyl, N_3 , NH_2 , NHBn or NH-acyl and the other is H, or V^1 and V^2 together are $=O$, $-O(CH_2)_pO-$ or $-S(CH_2)_pS-$;

W^1 is either O or S;

X^1 and X^2 are both H, or together are =O or =S;

Y is OR^5 or NR^6R^7 ;

Z is S or $-CH=CH-$;

R^1 , R^2 , R^3 and R^4 are independently selected from H, lower alkyl, lower alkyloxy, F, Cl and Br;

R^5 is selected from H and lower alkyl;

R^6 and R^7 are independently selected from H and lower alkyl, or together are $-(CH_2)_n-$;

R^8 is H or lower alkyl;

$n=3, 4, 5$ or 6 ;


p is 2 or 3 ; and

x is $0, 1$ or 2 .







The invention further comprises pharmaceutical compositions incorporating these vasopressin agonists, which compositions are particularly useful in the treatment of central diabetes insipidus, nocturnal enuresis and nocturia.

DESCRIPTION OF THE INVENTION

The present invention comprises N-benzylcarbamyl pyrrolidine derivatives defined by general formula 1.



In this formula, A represents a bicyclic or tricyclic azepine group according to one of the general formulae 2-7.



A¹, A⁴, A⁷ and A¹⁰ represent divalent groups selected from methylene (-CH₂-), oxygen (-O—) and substituted nitrogen (-NR⁸-). A², A³, A⁹, A¹¹, A¹³, A¹⁴ and A¹⁶ represent either a nitrogen atom (-N=) or a methine group (-CH=). A⁵ can represent a covalent bond, in which case A⁶ represents a sulphur atom (-S-) such that the ring that includes these two groups is a thiophene ring. Alternatively, A⁵ can represent a group -N=CH-, in which case A⁶ represents a covalent bond such that the ring that includes these two groups is a pyridine ring. A⁸ and A¹² represent either -NH- or a sulphur atom (-S-). A¹⁸ and A¹⁷ represent divalent groups. Both may be methylene groups (-CH₂-) or one is a methylene group and the other is selected from hydroxymethylene (-CH(OH)-), difluoromethylene (-CF₂-), oxygen (-O-), substituted nitrogen (-NR⁸-) and sulphur or oxidised sulphur (-S-, -SO-, or -SO₂-).

V¹ and V² may both be hydrogen, methoxy or fluorine, or one may be selected from bromine, chlorine, fluorine, hydroxy, lower alkoxy, benzyloxy, phenoxy, acyloxy, azido, amino, benzylamino and acylamido (Br, Cl, F, OH, O-lower alkyl, OBn, OPh, O-acyl, NH₂, NHBn and NH-acyl) provided that the other is hydrogen, or V¹ and V² together may represent an oxygen atom such that the fragment CV¹V² is a carbonyl group (C=O). V¹ and V² may also be an ethylene- or propylene-dioxy or -dithio chain (-O(CH₂)₂O-, -O(CH₂)₃O-, -S(CH₂)₂S-, -S(CH₂)₃S-) such that CV¹V² is a 1,3-dioxolane, 1,3-dioxane, 1,3-dithiolane or 1,3-dithiane ring.

W¹ is either an oxygen or a sulphur atom.

X¹ and X² may either both be hydrogen, or together they may represent an oxygen or sulphur atom such that the fragment CX¹X² is a carbonyl or thiocarbonyl group (C=O or C=S).

Y is either a group -OR⁶ or a group -NR⁶R⁷.

Z represents either a sulphur atom, such that the ring that includes it is a thiophene ring, or it represents a group $-\text{CH}=\text{CH}-$, such that the ring is a benzene ring.

R^1 , R^2 , R^3 and R^4 are each independently selected from hydrogen, lower alkyl groups, lower alkyloxy groups and the halogens fluorine, chlorine and bromine.

R^5 may be either a hydrogen atom or a lower alkyl group.

R^6 and R^7 may each independently be hydrogen atoms or lower alkyl groups, or together they may constitute a chain of between 3 and 6 methylene groups such that, together with the nitrogen atom to which they are attached, they form an azetidine, pyrrolidine, piperidine or perhydroazepine ring.

R^8 may be hydrogen or a lower alkyl group.

In the context of the present disclosure, the term "lower alkyl" is intended to include straight chain and branched alkyl groups and cycloalkyl groups of between 1 and 6 carbon atoms. For example, methyl, ethyl, isopropyl, *tert*-butyl, neopentyl and cyclohexyl are all within the scope of the term lower alkyl. The term "acyl" denotes lower alkyl carbonyl groups such as acetyl, pivaloyl, cyclopropylcarbonyl and the like. Formyl is also considered to be an acyl group.

Certain compounds of general formula 1 are capable of forming salts with acids or bases. For example, compounds containing one or more nitrogen atoms can form addition salts with mineral and organic acids such as hydrochloric acid, sulphuric acid, phosphoric acid, acetic acid, trifluoroacetic acid, methanesulphonic acid, citric acid and benzoic acid. Compounds containing acidic groups can form salts with bases. Examples of such salts include the sodium, potassium, calcium, triethylammonium and tetraethylammonium salts. Furthermore, compounds that have both acidic and basic groups can form internal salts (zwitterions). Insofar as these salts are pharmaceutically acceptable, they are included within the scope of the invention.

The compounds according to general formula 1 all have at least one stereogenic centre (a tetrahedral carbon atom bearing four different substituents) and so can exist as optical isomers such as enantiomers and diastereomers. Such isomers, and mixtures thereof, are all intended to be within the scope of the present invention.

In a preferred embodiment of the present invention, A is a group according to general formula 2. In another preferred embodiment of the present invention, A is a group

according to general formula 3. In another preferred embodiment of the present invention, A is a group according to general formula 4. In another preferred embodiment of the present invention, A is a group according to general formula 5. In another preferred embodiment of the present invention, A is a group according to general formula 6.

In another preferred embodiment of the present invention, A is a group according to general formula 7. In a more preferred embodiment, A is a tetrahydro-1-benzazepin-1-yl group, i.e. a group according to general formula 7 in which Z is $-\text{CH}=\text{CH}-$ and both A^{18} and A^{17} are methylene groups.


In another preferred embodiment, one of R^1 and R^2 is chlorine or a methyl group and the other is hydrogen, with both R^3 and R^4 also being hydrogen.

In another preferred embodiment, one of V^1 and V^2 is a methoxy or benzyloxy group and the other is hydrogen.

In yet another preferred embodiment, X^1 and X^2 together represent an oxygen atom and Y is $-\text{NR}^6\text{R}^7$.


Particularly preferred embodiments of the present invention are those that combine two or more of the above preferred features.

A still more preferred embodiment of the present invention is a compound according to general formula 8.




In general formula 8, W^1 , R^5 and R^6 are as defined above for 1. One of R^a and R^b is hydrogen and the other is either chlorine or a methyl group. R^c is either a methyl group or a benzyl group.

A yet more preferred embodiment is a compound of general formula 8A in which the stereochemistry is as shown.




Another preferred embodiment of the present invention is a compound according to general formula 1 in which V^1 and V^2 are both hydrogen. In a more preferred embodiment, X^1 and X^2 together are an oxygen atom and Y is NR^6R^7 . More preferred still is a compound according to general formula 9.



In general formula 9, W^1 , R^5 and R^6 are as defined above for 1. One of R^a and R^b is hydrogen and the other is either chlorine or a methyl group.

Even more preferred is a compound according to general formula 9A in which the stereochemistry is as shown.



Individual preferred compounds within the present invention include (but are not limited to) the following:

1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,

(4*R*)-4-hydroxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,

(4*R*)-1-(3-chloro-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide,

(4*R*)-1-(2-chloro-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide,

(4*R*)-4-benzyloxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,

(4*R*)-4-methoxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,

(4*R*)-4-methoxy-1-(3-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,


(4*R*)-1-(2-chloro-4-(5,6,7,8-tetrahydro-4*H*-thieno[3,2-*b*]azepin-4-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide,

(4*R*)-1-(4-(10,11-dihydro-5*H*-pyrrolo[2,1-*c*](1,4)benzodiazepin-10-ylcarbonyl)-2-methylbenzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide,

(4*R*)-1-(2-chloro-4-(10,11-dihydro-5*H*-pyrrolo[2,1-*c*](1,4)benzodiazepin-10-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide, and

(4*R*)-1-(4-(10,11-dihydro-5*H*-pyrrolo[2,1-*c*](1,4)benzodiazepin-10-ylcarbonyl)-2-methylbenzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylthioamide.

The compounds of the present invention can be prepared using methods generally known in the art. The compounds of general formula 1 can be considered to be composed of three linked fragments (A - C).




The three fragments will generally be prepared separately and then combined at a late stage in the synthesis. Some instances of the various groups ($R^1 - R^4$, V^1 , V^2 , X^1 , X^2 etc.) might be incompatible with this assembly and so will require the use of protecting groups. The use of protecting groups is well known in the art (see for example "Protective Groups in Organic Synthesis", T.W. Greene, Wiley-Interscience, 1981). Particular groups that may require protection are amines (protected as amides or carbamates), alcohols (protected as esters or


ethers) and carboxylic acids (protected as esters). For the purposes of this discussion, it will be assumed that such protecting groups as are necessary are in place.

The fragments A, B and C can be combined according to two strategies to give the compounds of formula 1. In the first, fragments A and B are linked to give a fragment corresponding to AB, which is then combined with fragment C. In the second, fragments B and C are linked to give a fragment corresponding to BC, which is then combined with fragment A. The chemistry involved in the condensation of fragment A with B, and that involved in the condensation of fragment B with fragment C, will be the same whichever strategy is followed. We have found that the first strategy is more flexible when working on a small scale and for preparing a selection of compounds. Nevertheless, it is possible that the second strategy would be advantageous for the preparation of a selected compound on a large scale.

Formation of fragment AB



Here, {A} and {B} represent part structures of the fragments A and B respectively. The formation of amides by the condensation of carboxylic acids with amines is well known. In general, the acid and the amine are mixed in an aprotic solvent such as dichloromethane or dimethylformamide in the presence of a condensing agent such as a carbodiimide (for example "water-soluble carbodiimide", which is *N*-ethyl-*N*-(3-dimethylaminopropyl)-carbodiimide) or a reactive phosphorus derivative (for example "BOP", which is (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate). The reaction may optionally be catalysed by a tertiary amine such as triethylamine or 4-dimethylaminopyridine. Alternatively, the carboxylic acid may be converted to a more reactive derivative such as the acid chloride. Such a derivative can then be reacted with the amine as described above but without the need for a condensing agent.


Formation of fragment BC

Formation of the urea or thiourea bond between fragments B and C can be most easily achieved by allowing the primary amine corresponding to fragment B to react with a derivative of carbonic acid such as phosgene (wherein LG above is chlorine) or carbonyldiimidazole (wherein LG is 1-imidazolyl) to form an intermediate carbamic acid derivative. When W¹ is sulphur rather than oxygen, thiophosgene or thiocarbonyldiimidazole is used. The reaction is conveniently carried out in an aprotic solvent such as dichloromethane or dimethylformamide in the presence of a tertiary amine such as triethylamine or *N,N*-diisopropylethylamine. After allowing sufficient time for the formation of the intermediate, the secondary amine corresponding to fragment C can be added to the reaction mixture. It is not necessary to isolate the intermediate carbamate derivative.

As a variation of this process, it is possible to reverse the order of addition of the amines corresponding to fragments B and C, such that the carbamate derivative is formed from the secondary amine, and the primary amine is added subsequently.

Overall then, the following intermediates are required for the synthesis of the compounds of the present invention


i) For fragment A



Fused azepines according to these general formulae can be prepared according to methods reported in the literature. See for example: Aranapakam *et al.*, Bioorg. Med. Chem. Lett. 1993, 1733; Artico *et al.*, Farmaco. Ed. Sci. 24, 1969, 276; Artico *et al.*, Farmaco. Ed. Sci. 32, 1977, 339; Chakrabarti *et al.*, J. Med. Chem. 23, 1980, 878; Chakrabarti *et al.*, J. Med. Chem. 23, 1980, 884; Chakrabarti *et al.*, J. Med. Chem. 32, 1989, 2573; Chimirri *et al.*, Heterocycles 36, 1993, 601; Grunewald *et al.*, J. Med. Chem. 39, 1996, 3539; Klunder *et al.*, J. Med. Chem. 35, 1992, 1887; Liegeois *et al.*, J. Med. Chem. 37, 1994, 519; Olagbemiro *et al.*, J. Het. Chem. 19, 1982, 1501; Wright *et al.*, J. Med. Chem. 23, 1980, 462; Yamamoto *et al.*, Tet. Lett. 24, 1983, 4711; and International patent application, publication number WO99/06403.

Some of them are items of commerce.


ii) For fragment B



Because the primary amine and the carboxylic acid groups are incompatible, they must be developed separately and protected. Substituted benzoic acids are well known, and the carboxylic acid is conveniently protected as its methyl ester. The primary amine can be elaborated from the corresponding nitrile (by reduction) or the alcohol (by displacement with

a nitrogen nucleophile). The best method will depend on the nature of the substituents R^1 - R^4 .

iii) For fragment C



Pyrrolidine derivatives of this type are prepared according to the methods described in the literature. See for example: Dugave *et al.*, Tet. Lett. 39, 1998, 1169; Petrillo *et al.*, J. Med. Chem. 31, 1988, 1148; and Smith *et al.*, J. Med. Chem. 31, 1988, 875.

Proline and hydroxyproline derivatives of defined stereochemistry are items of commerce and as such are convenient starting materials.

The present invention further comprises pharmaceutical compositions that include at least one compound according to the foregoing description as an active constituent. The composition may also include a second pharmacological agent such as a spasmolytic or a potassium channel blocker, these agents being known in the art to ameliorate bladder dysfunction. Preferably, the composition includes only one active constituent. The composition will include excipients selected from binding agents, bulking agents, dispersants, solvents, stabilising agents and the like, such excipients being generally known in the art.

The excipients used will depend on the intended nature of the formulation, which will, in turn, depend on the intended route of administration. Administration may be oral, transmucosal (such as sublingual, buccal, intranasal, vaginal and rectal), transdermal or by injection (such as subcutaneous, intramuscular and intravenous). Oral administration is generally preferred. For oral administration, the formulation will be a tablet or capsule. Other formulations include dry powders, solutions, suspensions, suppositories and the like.

In a further aspect, the present invention is a method of treating or controlling certain human physiological dysfunctions. This method comprises the administration to the person in need

of such treatment of an effective amount of a pharmaceutical composition, which composition contains a compound according to the foregoing description as an active constituent. The compounds act to reduce urine output, and so the method of the invention can be applied to all conditions in which elevated urine output is a contributory factor. The compounds also increase the production of the blood coagulation proteins known as Factor VIII and von Willebrand factor, and so the treatment of bleeding disorders can be undertaken.

In a preferred embodiment, the condition treated is diabetes insipidus. This is a condition caused by an inability of the body to produce and secrete physiologically active vasopressin, with the result that water re-uptake is greatly reduced and large volumes of urine are produced.

In another preferred embodiment, the condition treated is nocturnal enuresis. This is defined as bladder emptying while the individual is sleeping. It is a condition that mainly affects children and a number of factors may be involved in its etiology.

In another preferred embodiment, the condition treated is nocturia. This is defined as production of sufficient urine during the night to require the individual to wake and empty his (or her) bladder. Again, this condition may be the result of a number of factors.

In another preferred embodiment, the condition treated is incontinence. This condition is characterised, in part, by reduced bladder capacity and control such that involuntary urination occurs unless the bladder is emptied frequently. Incontinence has been divided into two conditions, stress incontinence and urge incontinence. A number of etiological factors are thought to be involved. Treatment according to the invention is particularly useful for delaying the need for bladder emptying ("voiding postponement") in order to allow the incontinent subject a dry period of a few hours (such as up to four hours). Such voiding postponement may also be useful for the non-incontinent population, for example for people obliged to remain in meetings for extended periods.

In another preferred embodiment, the condition treated is haemophilia A or von Willebrand's disease. These are conditions in which Factor VIII or von Willebrand factor production is reduced and the individual suffers from prolonged bleeding.

In another preferred embodiment, the composition is administered prior to surgery (including dental surgery) to increase the coagulability of the blood and so reduce peri-operative blood loss.

The administration of the compositions of the present invention will generally be under the control of a physician. The physician will determine the amount of composition to be administered and the dosing schedule, taking into account the patient's physical condition and the therapeutic goals. For an adult diabetes insipidus patient, a typical dose might be between 50mg and 1g of the active compound per day, taken as a single tablet or as up to four tablets throughout the day. For routes of administration other than the oral route, the amount of compound will be reduced, since non-oral routes tend to be more efficient in terms of delivering therapeutic agents into the systemic circulation. For the treatment of haemophilia A and von Willebrand's disease the amount of compound may need to be higher than for the treatment of diabetes insipidus.

The foregoing general description will now be further illustrated with a number of non-limiting examples.

EXAMPLES

Abbreviations.

The following abbreviations have been used.

Ac	Acetyl
AIBN	Azo-bis-(isobutyronitrile)
Bn	Benzyl
BOC	<i>tert</i> -Butyloxycarbonyl
(BOC) ₂ O	Di- <i>tert</i> -butyl dicarbonate
DMF	Dimethylformamide
Et	Ethyl
EtOAc	Ethyl acetate
IPA	Isopropanol
iPr	Isopropyl
M.S.	Mass spectrometry

Me	Methyl
NBS	N-Bromosuccinimide
pet. ether	petroleum ether, fraction boiling at 60-80°C
Ph	Phenyl
tBu	<i>tert</i> -Butyl
THF	Tetrahydrofuran
WSCDI	Water-soluble carbodiimide


Preparation of Intermediates.

Reagents corresponding to fragment A and C were commercially available or prepared according to the published procedures except where detailed in the specific Examples.

Reagents corresponding to fragment B were prepared as detailed below.

Example A.

4-(*tert*-Butyloxycarbonylaminoethyl)-3-chlorobenzoic acid




A1. Methyl 4-bromomethyl-3-chlorobenzoate


To a solution of methyl 3-chloro-4-methylbenzoate (5.0g, 27.1mmol) in carbon tetrachloride (50ml) were added NBS (5.8g, 32.0mmol) and AIBN (0.442g, 2.70mmol). The mixture was stirred at reflux for 18h. The mixture was allowed to cool to room temperature and then concentrated *in vacuo*. The residue was purified by flash chromatography on silica (eluant EtOAc:pet. ether 0:100 to 5:95); yield 5.96g (84%).

A2. 4-(*tert*-Butyloxycarbonylaminomethyl)-3-chlorobenzoic acid

To a saturated solution of ammonia in ethanol (170ml) was added methyl 4-bromomethyl-3-chlorobenzoate from Example A1 (5.5g, 20.9mmol). The mixture was stirred at room temperature for 1h and then concentrated *in vacuo*. The residue was triturated with diethyl ether and the resultant white crystals were filtered off and washed with more diethyl ether. To a solution of this solid in water (100ml) were added solutions of (BOC)₂O (5.0g, 23.0mmol) in dioxan (100ml) and sodium hydroxide (1.86g, 46.0mmol) in water (100ml). The mixture was stirred at room temperature for 18h and then concentrated *in vacuo*. The aqueous residue was acidified with citric acid and extracted with chloroform/IPA. The organic layer was washed with water, dried over MgSO₄, and concentrated *in vacuo* to give a white solid; yield 2.8g (67%).


Example B.4-Cyano-3-methylbenzoic acid

To a solution of 4-bromo-2-methylbenzonitrile (2.0g, 10.2mmol) in THF (100ml) at -78°C under a nitrogen atmosphere was added dropwise a 2.5M solution of *n*-butyl lithium (4.48ml, 11.2mmol). The mixture was stirred at -78°C for 1h and then poured onto solid carbon dioxide (5g) in THF (50ml). The mixture was allowed to warm to room temperature. Water was added (200ml) and the mixture was extracted with diethyl ether (3 times). The aqueous layer was acidified by addition of concentrated HCl and extracted with chloroform (3 times). The combined chloroform extracts were washed with water, dried over MgSO₄, and concentrated *in vacuo* to give a white solid; yield 1.2g (73%).

Example C.**4-Cyano-2-methylbenzoic acid**

4-Bromo-3-methylbenzonitrile (2.0g, 10.2mmol) was reacted following the method of Example B to give a yellow solid which was triturated with hexane and filtered off; yield 0.96g (59%).

Reagents corresponding to fragments A, B and C were combined to give the specific Examples as detailed below.

Example 1.**1-(2-Methyl-4-((2,3,4,5-tetrahydro-1-benzazepin-1-yl)carbonyl)benzylcarbamoyl)-L-proline-N,N-dimethylamide****1A. 2-Methyl-4-((2,3,4,5-tetrahydro-1H-benzo[b]azepine)-1-carbonyl)-benzonitrile.**

To a solution of 2,3,4,5-tetrahydro-1H-benzo[b]azepine (0.80g, 5.44mmol) in dichloromethane (50ml) were added 4-cyano-3-methylbenzoic acid (0.96g, 5.95mmol), triethylamine (0.60g, 5.95mmol), 4-(dimethylamino)pyridine (0.73g, 5.95mmol) and WSCDI (1.24g, 6.48mmol). The mixture was stirred at reflux for 18h, cooled and evaporated *in vacuo*. The residue was partitioned between EtOAc and 1M KHSO₄. The organic layer

was washed with saturated sodium bicarbonate solution and brine, dried over MgSO_4 , and concentrated *in vacuo*. The crude material was purified by flash chromatography on silica (eluant EtOAc:pet. ether 30:70); yield 1.10g (70%).

1B. 1-(4-(Aminomethyl)-3-methylbenzoyl)-2,3,4,5-tetrahydro-1H-benzof[b]azepine hydrochloride.


To a degassed solution of the cyanobenzazepine of Example 1A (1.10g, 3.79mmol) in methanol (50ml) were added concentrated hydrochloric acid (0.98ml, 11.3mmol) and 10% palladium on carbon (0.80g). Hydrogen gas was bubbled through the mixture for 5h at room temperature. The catalyst was removed by filtering through a pad of celite and the filtrate was evaporated; yield 1.23g (98%).

1C. 1-(2-Methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-N,N-dimethylamide

To a solution of the amine of Example 1B (0.10g, 0.302mmol) in DMF (10ml), under a nitrogen atmosphere, were added *N,N*-diisopropylethylamine (43mg, 0.332mmol) and carbonyl diimidazole (0.074g, 0.453mmol). The mixture was stirred at room temperature for 40 minutes. A solution of proline-*N,N*-dimethylamide (0.107g, 0.756mmol) in DMF (1ml) was added. The mixture was stirred at room temperature for a further 16 hr. The solvent was removed *in vacuo* and the crude material was purified by flash chromatography on silica (eluant methanol:dichloromethane 5:95); yield 0.115g (82%).

^1H NMR (CDCl_3): δ 1.35-1.55 (1H, m), 1.74-2.10 (3H, m), 2.11 (3H, s), 2.17-2.35 (1H, m), 2.60-2.82 (2H, m), 2.86 (3H, s), 2.90-3.14 (2H, m), 3.05 (3H, s), 3.26 (1H, dd, $J=14.9$ & 7.2Hz), 3.40-3.53 (1H, m), 3.64-3.84 (1H, m), 4.03-4.19 (1H, m), 4.29-4.42 (1H, m), 4.55-4.68 (1H, m), 4.74-4.81 (1H, m), 4.85-4.98 (1H, m), 6.58 (1H, d, $J=7.7\text{Hz}$), 6.75-6.89 (2H, m), 6.91-7.06 (3H, m), 7.16 (1H, d, $J=6.5\text{Hz}$), 7.93-8.03 (1H, m) ppm.

M.S.: calc $m/e=462.26$; found $[\text{M}+\text{H}]^+=463.2$

Example 2.**(4*R*)-4-Hydroxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzyl-carbamoyl)-L-proline-*N,N*-dimethylamide****2A. L-*trans*-4-Hydroxyproline-*N,N*-dimethylamide hydrochloride**

To a solution of BOC-hydroxyproline (2.99g, 13.89mmol) in dichloromethane (100ml) were added *N,N*-diisopropylethylamine (3.7ml, 21.24mmol), 4-(dimethylamino)pyridine (1.74g, 14.24mmol), dimethylamine hydrochloride (1.72g, 21.09mmol) and WSCDI (3.17g, 16.68mmol). The mixture was stirred at room temperature for 30hr. The mixture was diluted with dichloromethane (100ml) and washed with 0.3M KHSO₄, saturated sodium bicarbonate solution and brine, dried over MgSO₄, and concentrated *in vacuo* to give a colourless gum. This crude material was taken up in 4N HCl/dioxan (50ml) and stirred at room temperature for 1hr and then concentrated *in vacuo*. The residue was azeotroped with toluene and diethyl ether to give a white solid; yield 0.45g (17%).

2B. (4*R*)-4-Hydroxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzyl-carbamoyl)-L-proline-*N,N*-dimethylamide.

The amine of Example 1B (0.10g, 0.302mmol) was reacted with the amine of Example 2A (0.153mg, 0.785mmol) following the method of Example 1C. The product was purified by flash chromatography on silica (eluant chloroform:methanol:acetic acid 95:4:1); yield 0.95g (66%).


¹H NMR (CDCl₃): δ 1.65-1.80 (2H, m), 1.85-2.00 (3H, m), 2.05-2.25 (1H, m), 2.10 (3H, s), 2.80-3.10 (3H, m), 2.85 (3H, s), 3.00 (3H, s), 3.40-3.30 (1H, m), 3.45-3.55 (1H, m), 3.65-

3.95 (1H, m), 4.00–4.10 (1H, m), 4.30–4.55 (1H, m), 4.91 (1H, t, J=7.7Hz), 5.15–5.30 (1H, m), 6.10–6.20 (1H, m), 6.55–6.65 (1H, m), 6.85–7.50 (5H, m) ppm.


M.S.: calc m/e=478.26; found $[M+H]^+ = 479.2$

Examples 3 – 116.

The additional examples set out in the following Tables were prepared using analogous methods.




Example No	R/S	V ²	X	Y	$[M+H]^+$
3	<i>S</i>	H	H ₂	OMe	436.4
4	<i>R</i>	H	H ₂	OMe	436.2
5	<i>R/S</i>	OPh	O	OH	528.3
6	<i>R/S</i>	OPh	O	NMe ₂	555.3




Example No	R ¹	R ²	V ¹	V ²	X	Y	$[M+H]^+$
7	H	Me	H	H	O	OtBu	492.5
8	H	Me	H	H	O	OH	436.3
9	H	Me	H	OH	O	OMe	466.0

Example No	R ¹	R ²	V ¹	V ²	X	Y	[M+H] ⁺
10	H	Me	H	OAc	O	NMe ₂	521.0
11	H	Me		=O	O	NMe ₂	477.3
12	H	Me	H	OH	O	OEt	480.2
13	H	Me	H	OCOC ₃ H ₅	O	NMe ₂	547.3
14	H	Me	H	OMe	O	NMe ₂	493.5
15	H	Cl	H	H	O	NMe ₂	483.4
16	H	Me	H	H	S	NMe ₂	479.2
17	H	Me	H	H	O	NMeEt	477.2
18	H	OMe	H	H	O	NMe ₂	479.2
19	H	Me	H	OMe	O	OMe	480.2
20	H	Me	H	H	O	OIPr	478.2
21	H	Me	H	OH	O	OH	452.1
22	H	Me	H	OBn	O	OIPr	584.2
23	H	Me	H	OH	O	OiPr	494.1
24	H	Me	H	OBn	O	NMe ₂	569.2
25	Me	H	H	H	O	NMe ₂	463.2
26	H	Me	H	OMe	O	OH	466.2
27	Cl	H	H	H	O	NMe ₂	483.1
28	H	Et	H	H	O	NMe ₂	477.3
29	H	Cl	H	H	S	NMe ₂	499.2
30	H	Cl	H	OBn	O	NMe ₂	589.2
31	H	Cl	H	OH	O	NMe ₂	499.2
32	H	Me	H	OEt	O	NMe ₂	507.3
33	H	Me	Br	H	O	NMe ₂	541.1
34	H	Me	H	Cl	O	OMe	484.1
35	H	Me	F	F	O	NMe ₂	499.2
36	H	Me	H	Cl	O	OH	470.1
37	H	Me	H	N ₃	O	NMe ₂	504.3
38	H	Me	H	Cl	O	NMe ₂	497.2
39	H	Me	H	OtBu	O	NMe ₂	535.3
40	H	Me	Cl	H	O	NMe ₂	497.2
41	H	Me	H	OPh	O	OMe	542.3
42	H	Me	H	F	O	OMe	468.3

Example No	R ¹	R ²	V ¹	V ²	X	Y	[M+H] ⁺
43	H	Me	H	F	O	OH	454.4
44	H	Me	H	F	O	NMe ₂	481.3
45	H	Me	H	NHBn	O	NMe ₂	568.0
46	H	Me	OMe	OMe	O	OMe	510.3
47	H	Me	OMe	OMe	O	OH	496.2
48	H	Me	OMe	OMe	O	NMe ₂	523.3




Example No	R ²	V ²	X	[M+H] ⁺
49	Cl	H	O	489.1
50	Me	H	O	469.2
51	Me	OH	O	485.0
52	Cl	OMe	O	519.3
53	Me	OMe	O	499.3
54	Cl	OMe	S	535.1




Example No	R ³	V ²	W ¹	X	[M+H] ⁺
55	H	H	S	O	479.4
56	H	OH	S	O	495.0








Example No	R ³	V ²	W ¹	X	[M+H] ⁺
57	H	H	S	S	495.1
58	Me	H	O	O	477.2
59	H	OBn	S	O	585.2
60	H	OBn	O	S	585.0







Example No	A ¹⁶	A ¹⁷	R ²	V ²	[M+H] ⁺
61	NEt	CH ₂	Me	OMe	522.4
62	NH	CH ₂	Me	OMe	494.3
63	CH ₂	NiPr	Me	OMe	536.4
64	CH ₂	NH	Me	OMe	494.5
65	O	CH ₂	Cl	OMe	515.2
66	CH(OH)	CH ₂	Me	H	479.2




Example No	A	V ²	[M+H] ⁺
67		H	518.0

Example No	A	V ²	[M+H] ⁺
68		H	532.2
69		H	527.0
70		H	516.1
71		H	515.0
72		H	514.6
73		H	513.7
74		H	502.1


Example No	A	V ²	[M+H] ⁺
75		H	500.7
76		OH	547.9
77		OH	517.6
78		OH	546.3
79		H	517.2
80		OBn	619.2
81		OMe	543.4
82		OMe	544.3


Example No	A	V ²	[M+H] ⁺
83		OMe	549.2
84		OMe	548.2
85		OMe	562.1
86		OMe	590.2




Example No	V ¹	V ²	X	Y	[M+H] ⁺
87	H	H	S	NMe ₂	516.2
88	H	OBn	O	NMe ₂	606.3
89	H	OH	O	NMe ₂	507.3
90	H	OMe	O	NMe ₂	530.3
91	-OCH ₂ CH ₂ O-		O	OMe	545.3
92	OMe	OMe	O	OMe	547.3

Example No	V ¹	V ²	X	Y	[M+H] ⁺
93	-OCH ₂ CH ₂ O-		O	NMe ₂	558.3
94	-SCH ₂ CH ₂ S-		O	NMe ₂	590.2



Example No	R ²	V ¹	V ²	X	Y	[M+H] ⁺
95	Me	H	OH	O	NMe ₂	516.1
96	Me	H	H	S	NMe ₂	516.2
97	Me	H	OMe	O	NMe ₂	530.4
98	Me	-OCH ₂ CH ₂ O-		O	OMe	545.3
99	Me	-OCH ₂ CH ₂ O-		O	OH	531.3
100	Me	-OCH ₂ CH ₂ O-		O	NMe ₂	558.3
101	Cl	H	H	O	NMe ₂	551.5
102	Me	H	OMe	O	NEt ₂	558.3
103	Me	H	OMe	O		570.3
104	Me	H	OMe	S	NMe ₂	546.2



Example No	A ¹⁰	R ²	V ²	X	[M+H] ⁺
105	O	Me	H	O	519.3
106	NMe	Me	H	O	532.3
107	NMe	Me	OH	O	548.1
108	NMe	Me	OBn	O	638.2
109	NMe	Me	OMe	O	562.3
110	O	Me	OMe	O	549.2
111	NMe	Me	Cl	O	566.2
112	NMe	Me	OMe	S	578.2
113	O	Cl	OMe	O	569.1
114	O	Me	OMe	S	565.2
115	O	Cl	OMe	S	585.1
116	NH	Me	OMe	O	548.2

Example 117.

In Vitro Biological Characterisation

The compounds of the invention are selective agonists at the V₂ receptor. In standard radio-ligand displacement assays, the compounds all give K_i values below 10 μM for the V₂ receptor.

Example 118.**In Vivo Biological Characterisation**

The Brattleboro rat is a recognised model for vasopressin deficiency (for a review see FD Grant, "Genetic models of vasopressin deficiency", *Exp. Physiol.* 85, 203S-209S, 2000). The animals do not secrete vasopressin and consequently produce large volumes of dilute urine. Compounds of the invention were administered to Brattleboro rats (0.1-10mg/kg p.o. in methylcellulose). Urine was collected hourly and volumes were compared with control animals. Animals had free access to food and water throughout the experiment. Representative results are given in the Table. Results for Desmopressin are given for comparison.

Compound of Example	Dose	% Inhibition of urine output (at 1 hour)
1	1mg/kg	82
14	1mg/kg	84
52	1mg/kg	90
54	1mg/kg	68
85	1mg/kg	63
90	1mg/kg	60
101	1mg/kg	74
104	1mg/kg	81
109	1mg/kg	73
110	1mg/kg	80
112	1mg/kg	75
114	1mg/kg	85
115	1mg/kg	88
Desmopressin	0.1mg/kg	37
	1mg/kg	100
	10mg/kg	100

Example 119.**Pharmaceutical composition for tablet**

Tablets containing 100mg of the compound of Example 1 as the active agent are prepared from the following:

Compound of Example 1	200.0g
Corn starch	71.0g
Hydroxypropylcellulose	18.0g
Carboxymethylcellulose calcium	13.0g
Magnesium stearate	3.0g
Lactose	195.0g
<i>Total</i>	<i>500.0g</i>

The materials are blended and then pressed to give 2000 tablets of 250mg, each containing 100mg of the compound of Example 5.


The foregoing Examples demonstrate that compounds within the scope of the invention are readily prepared using standard chemical techniques, and that these compounds have the biological properties that would be expected of V_2 receptor agonists. In particular, the compounds are potent antidiuretics in an animal model of vasopressin deficiency. Thus it is clear that they may be useful in the treatment of human diseases that are currently treatable with Desmopressin, such as central diabetes insipidus, nocturnal enuresis and nocturia. It has further been suggested that antidiuretics such as Desmopressin may be useful in certain types of urinary incontinence. These arguments would also extend to the compounds of the present invention.

Desmopressin is also used in the treatment of certain coagulation disorders. There is good evidence to suggest that this action is also mediated through the V_2 receptor (see for example JE Kaufmann *et al.*, "Vasopressin-Induced von Willebrand factor secretion from endothelial cells involves V_2 receptors and cAMP", *J. Clin. Invest.* **106**, 107-116, 2000; A Bernat *et al.*, " V_2 receptor antagonism of DDAVP-induced release of hemostasis factors in conscious dogs", *J. Pharmacol. Exp. Ther.* **282**, 597-602, 1997), and hence it would be expected that the compounds of the present invention should be useful pro-coagulants.

The scope of the present invention is further defined in the following Claims.







CLAIMS

1. A compound according to general formula 1, or a pharmaceutically acceptable salt thereof,



wherein:

A is a bicyclic or tricyclic azepine derivative selected from general formulae 2 to 7



A¹, A⁴, A⁷ and A¹⁰ are each independently selected from CH₂, O and NR⁸;

A², A³, A⁹, A¹¹, A¹³, A¹⁴ and A¹⁵ are each independently selected from CH and N;

either A⁵ is a covalent bond and A⁶ is S, or A⁵ is N=CH and A⁶ is a covalent bond;

A⁸ and A¹² are each independently selected from NH and S;

A¹⁶ and A¹⁷ are both CH₂, or one of A¹⁶ and A¹⁷ is CH₂ and the other is selected from CH(OH), CF₂, O, SO_x, and NR⁸,

V¹ and V² are both H, OMe or F, or one of V¹ and V² is OH, OMe, OBn, OPh, O-acyl, Br, Cl, F, N₃, NH₂, NHBn or NH-acyl and the other is H, or V¹ and V² together are =O, -S(CH₂)_pS- or -O(CH₂)_pO-;

W¹ is either O or S;

X¹ and X² are both H, or together are =O or =S;

Y is OR⁵ or NR⁶R⁷;

Z is S or -CH=CH-;

R¹, R², R³ and R⁴ are independently selected from H, lower alkyl, lower alkyloxy, F, Cl and Br;

R⁵ is selected from H and lower alkyl;

R⁶ and R⁷ are independently selected from H and lower alkyl, or together are -(CH₂)_n-;

R⁸ is H or lower alkyl;


n=3, 4, 5 or 6;

p=2 or 3; and

x is 0, 1 or 2.

2. A compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein A is a group according to general formula 2.

3. A compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein A is a group according to general formula 3.
4. A compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein A is a group according to general formula 4.
5. A compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein A is a group according to general formula 5.
6. A compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein A is a group according to general formula 6.
7. A compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein A is a group according to general formula 7.
8. A compound according to Claim 1 or Claim 7, or a pharmaceutically acceptable salt thereof, wherein A is a group according to general formula 7, Z is $-\text{CH}=\text{CH}-$ and both A^{16} and A^{17} are $-\text{CH}_2-$.
9. A compound according to any of Claims 1 to 8, or a pharmaceutically acceptable salt thereof, wherein one of R^1 and R^2 is Cl or Me and the other is H, and both R^3 and R^4 are H.
10. A compound according to any of Claims 1 to 9, or a pharmaceutically acceptable salt thereof, wherein one of V^1 and V^2 is OMe or OBn and the other is H.
11. A compound according to any of Claims 1 to 10, or a pharmaceutically acceptable salt thereof, wherein X^1 and X^2 together are $=\text{O}$ and Y is NR^6R^7 .
12. A compound according to any of Claims 1 or 7 to 11, or a pharmaceutically acceptable salt thereof, which is



wherein

W^1 is either O or S;


one of R^a and R^b is Cl or methyl and the other is H;

R^c is methyl or benzyl;

R^6 and R^7 are independently selected from H and lower alkyl, or together are $-(CH_2)_n-$; and

n is 3, 4, 5 or 6.

13. A compound according to any of Claims 1 or 7 to 12, or a pharmaceutically acceptable salt thereof, which is



wherein

W^1 is either O or S;

one of R^a and R^b is Cl or methyl and the other is H;


R^c is methyl or benzyl; and

R^6 and R^7 are independently selected from H and lower alkyl, or together are $-(CH_2)_n-$; and

n is 3, 4, 5 or 6.

14. A compound according to any of Claims 1 to 9, or a pharmaceutically acceptable salt thereof, wherein V^1 and V^2 are both H.
15. A compound according to Claim 14, or a pharmaceutically acceptable salt thereof, wherein X^1 and X^2 together are $=O$ and Y is NR^6R^7 .

16. A compound according to any of Claims 1 or 7 to 9 or 14 or 15, or a pharmaceutically acceptable salt thereof, which is



wherein


W¹ is either O or S;

one of R^a and R^b is Cl or methyl and the other is H;

R⁶ and R⁷ are independently selected from H and lower alkyl, or together are -(CH₂)_n;
and

n is 3, 4, 5 or 6.

17. A compound according to any of Claims 1 or 7 to 9 or 14 to 16, or a pharmaceutically acceptable salt thereof, which is



wherein

W^1 is either O or S;

one of R^a and R^b is Cl or methyl and the other is H;

R^6 and R^7 are independently selected from H and lower alkyl, or together are $-(CH_2)_n-$; and

n is 3, 4, 5 or 6.

18. A compound according to any preceding Claim which is selected from

1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,

(4*R*)-4-hydroxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,

(4*R*)-1-(3-chloro-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide,

(4*R*)-1-(2-chloro-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide,

(4*R*)-4-benzyloxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,

(4*R*)-4-methoxy-1-(2-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,

(4*R*)-4-methoxy-1-(3-methyl-4-(2,3,4,5-tetrahydro-1-benzazepin-1-ylcarbonyl)benzylcarbamoyl)-L-proline-*N,N*-dimethylamide,

(4*R*)-1-(2-chloro-4-(5,6,7,8-tetrahydro-4*H*-thieno[3,2-*b*]azepin-4-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide,

(4*R*)-1-(4-(10,11-dihydro-5*H*-pyrrolo[2,1-*c*](1,4)benzodiazepin-10-yl carbonyl)-2-methylbenzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide,

(4*R*)-1-(2-chloro-4-(10,11-Dihydro-5*H*-pyrrolo[2,1-*c*](1,4)benzodiazepin-10-ylcarbonyl)benzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylamide, and

(4*R*)-1-(4-(10,11-dihydro-5*H*-pyrrolo[2,1-*c*](1,4)benzodiazepin-10-ylcarbonyl)-2-methylbenzylcarbamoyl)-4-methoxy-L-proline-*N,N*-dimethylthioamide

or a pharmaceutically acceptable salt thereof.

19. A use for a compound according to any of Claims 1 to 18, or a pharmaceutically acceptable salt thereof, which is as a component of a pharmaceutical composition.
20. A use for a compound according to any of Claims 1 to 18, or a pharmaceutically acceptable salt thereof, which is as a therapeutic agent for the treatment of nocturnal enuresis, nocturia, polyuria resulting from central diabetes insipidus, urinary incontinence and bleeding disorders.
21. A pharmaceutical composition which contains active agent selected from compounds and pharmaceutically acceptable salts according to any of Claims 1 to 18.

22. A pharmaceutical composition according to Claim 21, which composition is for the treatment of polyuria.
23. A pharmaceutical composition according to Claim 21, which composition is for the control of urinary incontinence.
24. A pharmaceutical composition according to Claim 21, which composition is for voiding postponement.
25. A pharmaceutical composition according to Claim 21, which composition is for the treatment of bleeding disorders.
26. A method of treatment of nocturnal enuresis, nocturia and central diabetes insipidus, which method comprises the administration to a person in need of such treatment of an effective amount of a composition according to Claim 21.
27. A method for the control of urinary incontinence, which method comprises the administration to a person in need of such treatment of an effective amount of a composition according to Claim 21.
28. A method of treatment according to Claim 27, wherein the treatment results in voiding postponement.
29. A method of treatment of bleeding disorders, which method comprises the administration to a person in need of such treatment of an effective amount of a composition according to Claim 21.

INTERNATIONAL SEARCH REPORT

Inter 1st Application No

PCT/GB 01/00023

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D403/12 C07D495/04 C07D471/04 C07D487/04 C07D498/04
 A61K31/55 A61P13/00 //(C07D495/04,223:00,333:00),
 (C07D495/04,243:00,333:00),(C07D471/04,221:00,243:00),

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 753 644 A (H. OGAWA ET AL.) 19 May 1998 (1998-05-19) claims 1-7 ---	1-29
A	WO 99 06403 A (AMERICAN HOME PRODUCTS CORPORATION) 11 February 1999 (1999-02-11) cited in the application claims 1-14 ---	1-29
A	WO 99 06409 A (AMERICAN HOME PRODUCTS CORPORATION) 11 February 1999 (1999-02-11) cited in the application claims 1-20 ---	1-29
A	WO 94 20473 A (YAMANOUCHI PHARMACEUTICAL CO., LTD.) 15 September 1994 (1994-09-15) claims 1-21 ---	1-29
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

21 March 2001

Date of mailing of the international search report

27/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

Herz, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/00023

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 (C07D487/04, 231:00, 243:00), (C07D487/04, 209:00, 255:00),
(C07D487/04, 243:00, 249:00), (C07D487/04, 209:00, 243:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 514 667 A (OTSUKA PHARMACEUTICAL CO., LTD.) 25 November 1992 (1992-11-25) claims 1-31 ---	1-29
A	WO 94 12476 A (YAMANOUCHI PHARMACEUTICAL CO., LTD.) 9 June 1994 (1994-06-09) claims 1-3 ---	1-29
A	PATENT ABSTRACTS OF JAPAN vol. 1999, no. 08, 30 June 1999 (1999-06-30) & JP 11 060488 A (OTSUKA PHARMACEUTICAL CO., LTD.), 2 March 1999 (1999-03-02) abstract --- -/--	1-29



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

21 March 2001

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Herz, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/00023

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>PATENT ABSTRACTS OF JAPAN vol. 018, no. 140, 8 March 1994 (1994-03-08) & JP 05 320135 A (YAMANOUCHI PHARMACEUTICAL CO., LTD.), 3 December 1993 (1993-12-03) abstract</p> <p>-----</p>	1-29

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 01/00023

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5753644	A	19-05-1998	AU 663628 B	12-10-1995
			AU 5161493 A	09-05-1994
			CA 2124696 A	28-04-1994
			CN 1098716 A, B	15-02-1995
			CN 1183277 A	03-06-1998
			EP 0620003 A	19-10-1994
			WO 9408582 A	28-04-1994
			JP 2913082 B	28-06-1999
			JP 6211800 A	02-08-1994
			MX 9306425 A	31-05-1994
			US 5622947 A	22-04-1997
WO 9906403	A	11-02-1999	AU 8592598 A	22-02-1999
			BR 9811559 A	12-09-2000
			CN 1272110 T	01-11-2000
			EP 1000059 A	17-05-2000
			NO 20000241 A	06-03-2000
WO 9906409	A	11-02-1999	AU 8663398 A	22-02-1999
			BR 9811585 A	26-09-2000
			CN 1272111 T	01-11-2000
			EP 1000062 A	17-05-2000
			HU 0002480 A	28-11-2000
			NO 20000242 A	06-03-2000
WO 9420473	A	15-09-1994	AU 6219994 A	26-09-1994
EP 514667	A	25-11-1992	AU 646334 B	17-02-1994
			AU 1498492 A	22-10-1992
			CA 2066104 A	20-10-1992
			CN 1066653 A, B	02-12-1992
			DE 69203955 D	14-09-1995
			DE 69203955 T	15-02-1996
			DK 514667 T	25-09-1995
			ES 2078576 T	16-12-1995
			JP 2916536 B	05-07-1999
			JP 5132466 A	28-05-1993
			KR 196485 B	15-06-1999
			MX 9201819 A	01-12-1992
			US 5244898 A	14-09-1993
WO 9412476	A	09-06-1994	AU 5533894 A	22-06-1994
JP 11060488	A	02-03-1999	JP 2926335 B	28-07-1999
JP 05320135	A	03-12-1993	NONE	